


Stratigraphic Unit	DESCRIPTION OF UNITS
Qa	Alluvium —Moderately to well-sorted clay, silt, sand, and gravel deposited by streams. Thickness ranges from 0 to 50 feet (DeWitt and others, 1989). A local aquifer where saturated.
Qc	Colluvium —Poorly to well-sorted, massive to stratified, clay-free to boulder rubble and talus produced by mass wasting. Thickness ranges from 0 to 50 feet (DeWitt and others, 1989). The hydrogeologic character of colluvial material varies considerably depending on the degree of sorting and primary type of material. Generally not an aquifer even if saturated.
Qp	Gravel deposits —Moderately sorted, heterogeneous, generally stratified, clay, silt, sand, and well-sorted gravel of paleochannels, pediments, and stream terraces along former flood plains. Three Quaternary terraces are identifiable in the northern portion of the study area and between six and eight terraces in the southern portion (Kempson and Lamy, 1994). Thickness ranges from 0 to 60 feet thick (DeWitt and others, 1989). A local aquifer where saturated.
Qw	Wind-blown deposits —Moderately to well-sorted, poorly to well-stratified, well-sorted, commonly calcareous, and frosted silt and fine to medium-grained sand. Thickness ranges from 0 to 50 feet thick (DeWitt and others, 1989). A local aquifer where saturated.
Tu	Gravel deposits —Heterogeneous gravel deposits derived primarily from igneous and metamorphic rocks of the central Black Hills. Clast size is dominantly pebble or cobble, but ranges from clay to boulder. Thickness ranges from 0 to 30 feet (J.F. Sawyer and J.E. Martin, South Dakota Geological Survey, written commun., 1997). A local aquifer where saturated.
Tr	White River Group —Consists of the Brule and Chadron Formations. Sandstone, claystone, and siltstone with channel fillings and limestone lenses (Rahn, 1985). Thickness ranges from 0 to 100 feet (DeWitt and others, 1989). A minor aquifer where saturated.
Tu	Undifferentiated shallow intrusive igneous rocks —Includes rhyolite, latite, trachyte, and phonolite (see DeWitt and others, 1989, for a more thorough description). The hydrogeologic characteristics of these rocks vary with the degree of fracturing.
Kps	Pierre Shale to Skull Creek Shale, undifferentiated —Confining unit of shale, limestone, and sandstone containing the following formations listed with their thickness in feet: Pierre Shale, 1,260-2,700; Niobrara Formation, 80-300; Carlile Shale, 350-750; Greenhorn Formation, 225-300; Belle Fourche Shale, 150-800; Osage Shale, 125-200; Newcastle Sandstone, 0-150; and Skull Creek Shale, 150-270 (modified from DeWitt and others, 1989). Where present, the Newcastle Sandstone is an aquifer if saturated.
Kk	Inyan Kara Group —Sandstone and other classic rocks of the Fall River Formation and Lakota Formation. The Fall River Formation is 100-200 feet thick consisting of brown to reddish-brown fine-grained sandstone, thin-bedded at the top and massive at the bottom. The Lakota Formation is 35-700 feet thick consisting of yellow, brown, and reddish-brown massive to thin-bedded sandstone, pebbles and conglomerate, siltstone, and claystone with local limestone, coal, and fossiliferous layers (DeWitt and others, 1989). A major regional aquifer.
Ju	Morrison Formation to Gypsum Spring Formation, undifferentiated —Semi-confining unit consisting of interbedded shale, sandstone, and gypsum of the following formations listed with their thickness in feet: Morrison Formation, 0-200; Utopia Sandstone, 0-225; Sundance Formation, 250-450; and Gypsum Spring Formation, 0-45 (modified from DeWitt and others, 1989). Sandstones of the Sundance Formation are locally productive aquifers where saturated (Kyllonen and Peter, 1987).
Tp	Spearfish Formation —Red siltstone interbedded with friable, red sandstone and siltstone, and sparse limestone layers. Lower portion contains massive gypsum (Robinson and others, 1964). Thickness ranges from 275 to 800 feet (Gries and Martin, 1995).
Pnk	Minnekahta Limestone —Fine-grained, purple to gray laminated limestone. Thickness ranges from 25 to 65 feet (modified from DeWitt and others, 1989). Unit is locally fractured and brecciated due to solution collapse (Gries and Martin, 1995). A major aquifer in the study area.
Op	Opeche Shale —Red siltstone and sandy shale, with local gypsum and anhydrite near the top. Thickness ranges from 25 to 150 feet (DeWitt and others, 1989).
Pph	Minnekahta Formation —Variably colored but generally yellow to red cross-stratified sandstone, limestone, dolomite, and shale of the Minnekahta Formation. Thickness ranges from 375 to 1,175 feet (modified from DeWitt and others, 1989), most commonly between 400 to 750 feet (Gries and Martin, 1995). Anhydrite is common at depth, generally within the upper 200 feet of the formation. Where anhydrite has dissolved, collapse-formed secondary permeability creates the most productive aquifer. Interbedded limestone and shale in the lower part of the formation form a confining zone (Kyllonen and Peter, 1987). A major regional aquifer.
MDne	Madison (Pahasapa) Limestone and Englewood Formation —Gray to buff and lavender limestone that is locally dolomitic. The Madison Limestone is 250 to 1,600 feet thick, and the Englewood Formation is 30 to 40 feet thick (modified from DeWitt and others, 1989). Generally massive, upper third is karstic with caves, solution collapse and enlarged conduits resulting in extensive secondary permeability and creating the potentially most productive aquifer in the Black Hills (Kyllonen and Peter, 1987). The lower portion of the Madison Limestone and the Englewood Formation form a lower confining zone (Greene, 1993). A major regional aquifer.
OCd	Whitewood Formation and Wilmot Formation —Undifferentiated semi-confining unit consisting of limestone and dolomite of the Whitewood Formation, thickness ranges from 0 to 235 feet, and shale with interbedded siltstone of the Wilmot Formation, thickness ranges from 0 to 150 feet (modified from DeWitt and others, 1989). The unit is found in the northern and western portion of the study area.
OCg	Deadwood Formation —Brown to light-gray glauconitic sandstone, shale, limestone, and local calcarenite. Thickness ranges from 25 to 500 feet (modified from DeWitt and others, 1989). A major aquifer in the study area.
Xp	Harney Peak Granite —Pink and tan coarse-grained and pegmatitic muscovite granite. Characterized geochronologically from other granites in the area by high concentrations of boron, beryllium, lithium, and strontium (DeWitt and others, 1989).
Xu	Undifferentiated metamorphosed phyllite and schist —Locally carbonaceous and lufaceous. Geochemical and geophysical signatures vary with the tuffaceous zones having an anomalously high copper concentration and a magnetic nature (DeWitt and others, 1989).
XWu	Undifferentiated igneous rocks —Geochemical signature and geophysical nature vary depending on position (see DeWitt and others, 1989 for additional information).
XWg	Metamorphosed graywacke —Primarily a medium- to dark-gray siliceous mica schist and impure quartzite (DeWitt and others, 1989).
XWv	Undifferentiated metamorphosed sedimentary deposits —Includes conglomerate, quartz sandstone, siltstone, and dolomite gneiss. Parts are characterized geochronologically by anomalously high uranium, chromium, and gold concentrations and other parts by anomalously high silver and arsenic concentration. The latter zones are magnetic in nature (DeWitt and others, 1989).

NOTE: Some thickness ranges given above are from the sources listed and from data file at the U.S. Geological Survey, Rapid City, South Dakota.

Planimetric base from U.S. Geological Survey digital data 1:100,000, Devils Tower and Sandstone, 1979; Belle Fourche, 1980; Rapid City, 1977. Topographic base modified from U.S. Geological Survey digital data, 1:24,000, from maps dated 1950-54. Universal Transverse Mercator projection. Zone 13. North American Horizontal Datum 1927.

EXPLANATION

- Fault (Dashed where approximated, dotted where concealed. Bar and tail on downthrown side.)
- Anticline (Showing trace of axial plane and direction of plunge. Dashed where approximated, dotted where concealed.)
- Syncline (Showing trace of axial plane and direction of plunge. Dashed where approximated, dotted where concealed.)
- Monocline (Showing trace of axial plane. Dashed where approximated, dotted where concealed.)
- Dome (Symbol size approximately proportional to size of dome. Dome asymmetry indicated by arrow length.)



Distribution of Hydrogeologic Units in the Black Hills Area, South Dakota
By Michael L. Strobel, Gregory J. Jarrell, J. Foster Sawyer, John R. Schleicher, and Mark D. Fahrenbach
1999