

A model for radionuclides in glacial and bedrock aquifers of the United States

Joseph D. Ayotte¹, Sarah M. Flanagan¹, Laura Hayes¹, and William S. Morrow² – U.S. Geological Survey, ¹New Hampshire – Vermont Water Science Center, ²Illinois Water Science Center

Introduction

Uranium and ²²²Rn are naturally occurring radionuclides found in ground water throughout the United States and can be associated with adverse health outcomes. This study uses a geologic model to explain the variation in uranium and ²²²Rn concentrations in three of the principal aquifers identified by the USGS's National Water Quality Assessment Program (NAWQA) (Figure 1).

Data from over 1,400 domestic, public-supply, and monitoring wells sampled by the NAWQA program from 1991-2003 were analyzed. Because of the size and complexity of the Glacial aquifers, we developed a model to differentiate sub aquifers based on source materials, glacial lobe delineation, and directions of Wisconsinian ice flow from Canada.

Figure 1. Locations of NAWQA Regional Assessments in Principal Aquifers



Incorporating a geologic model




Figure 2. Geology of the upper Midwest, United States

Figure 3. Wisconsinian glacial lobes and ice directions

To analyze regional variation of radionuclides in ground water in the glaciated U.S., the glacial principal aquifer was divided into sub-aquifers. Different classifications of the glacial aquifers are available (Meinzer, 1923; Heath, 1984; USGS, 2003) but are mostly based on age and stratigraphy. Other studies have used generalized geologic regions or political boundaries (Hess and others, 1985; Michael, 1987; Focazio and others, 2001; Longtin, 1988).

Maps of generalized bedrock geology (Fig. 2), Wisconsinian glacial lobes (Fig. 3), their advance directions, and their composition, were used to delineate important sub-aquifers within the glaciated terrain of the U.S. Analysis of the bedrock lithology and the glacial lobe data identified groups that might relate to radionuclide occurrence in ground water (King and Beikman, 1974; Schumann, 1993).

Incorporating terrestrial gamma surveys




Figure 4. Equivalent Thorium (ppm) and concentrations of uranium in wells

In addition to the glacial and bedrock geology, national data on the equivalent uranium (eU) and thorium (eTh) content of near-surface sediments from terrestrial gamma surveys (Duval, 1990) was used to help define areas of potential high concentrations of radionuclides in groundwater (Fig. 4). The 75th percentiles of uranium and ²²²Rn concentrations in ground water for each NAWQA study network are plotted against the associated 75th percentile of eU and eTh from the gamma survey. Above 2.0 ppm eU and 6 ppm eTh, ground water ²²²Rn concentrations generally increase (Fig. 5a, b). Above 5 ppm eTh, uranium concentrations increase but are highly variable (Fig. 5d).




Figure 5a-d. Relation of the 75th percentile concentration of ground water (a, b) ²²²Rn and (c, d) uranium to the 75th percentile eU and eTh in near-surface sediments.

A framework for regional analyses

The Principal Aquifer and sub-aquifers are shown on the maps below (Fig. 6a-b). The maps also show locations of wells used in this study and concentrations of uranium (Fig. 6a) and ²²²Rn (Fig. 6b).

WELL	Principal Aquifer	Sub-aquifer
PMG	PMG	PMG
CPG	CPG	CPG
WCS	WCS	WCS
NEC	NEC	NEC
ECG	ECG	ECG
PWG	PWG	PWG
COS	COS	COS
NEG	NEG	NEG
NEC	NEC	NEC




Figure 6a-b. Principal Aquifer and sub-aquifer groups with (a) uranium and (b) ²²²Rn in ground water. New England Crystalline underlies Northeastern Glacial aquifer. The Cambrian – Ordovician Sandstone aquifer underlies the Midwest Glacial aquifer. The Columbia Plateau Glacial aquifers are outside the extent of Pleistocene glaciation.

Source-water guideline exceedences

Table 2. Percent of drinking water wells and monitoring wells with uranium and ²²² Rn greater than the USEPA Maximum Contaminant Level (MCL or AMCL), by aquifer category and well type, ... no data									
Principal Aquifer or sub aquifer	Drinking water wells:				Monitoring wells:				
	Uranium		²²² Rn		Uranium		²²² Rn		
	N	Percent > 30 µg/L	N	Percent > 4,000 pCi/L	N	Percent > 30 µg/L	N	Percent > 300 pCi/L	
PMG	31	0	57	0	46	5	0	17	0
CPG	—	—	—	—	89	46	9	39	0
WCS	52	12	35	0	40	116	11	41	0
NEC	76	0	74	0	34	93	0	31	0
ECG	70	0	123	0	45	209	0	209	0
PWG	59	0	88	0	41	8	0	83	0
COS	73	0	74	0	66	—	—	—	—
NEG	67	0	61	2	32	55	0	29	0
NEC	113	4	107	30	94	—	—	—	—

Correlations with uranium and ²²²Rn

Spearman correlation coefficients (rho) between uranium and other water chemistry data by aquifer and sub aquifer. Analysis of correlations may help identify processes affecting concentrations of uranium in ground water (Table 3).

Table 3. Spearman correlation coefficients for Uranium by principal aquifer and sub aquifer; --, no data; bold p < 0.05.													
Aquifer	ANC	B	Ca	HCO3	Li	Mn	Ni	Se	SD4	Sr	TDS	TH	alpha
PMG	—	—	—	—	—	—	—	—	—	—	—	—	—
CPG	0.68	0.42	0.31	0.77	0.47	0.22	0.31	0.37	0.51	0.62	0.80	0.16	—
WCS	0.40	0.36	0.37	0.37	0.43	-0.31	0.80	0.37	0.32	0.38	0.06	—	—
NEC	0.11	0.05	0.19	0.24	0.20	0.19	0.42	0.19	0.38	0.44	0.17	0.16	0.08
ECG	0.35	0.16	0.40	0.29	0.11	0.48	0.45	0.16	0.41	0.23	0.36	0.16	—
PWG	-0.19	-0.45	0.47	—	—	-0.34	0.45	0.12	0.58	0.61	0.29	0.41	—
NEG	0.31	-0.03	0.29	0.30	0.16	0.31	-0.01	—	0.37	0.22	0.19	0.01	0.54
COS	0.48	-0.14	0.42	—	-0.22	0.36	0.06	0.32	0.02	0.27	0.09	—	—
NEC	0.16	-0.05	0.25	-0.03	0.24	-0.16	-0.01	-0.13	-0.03	0.10	0.14	0.80	0.72

New England Crystalline aquifers

In New England (NEC), uranium correlates with gross alpha (0.72) and ²²²Rn (0.60) (Fig. 7a-c). This unusual result was also reported in Finland (Vesterbacka, 2005), in similar bedrock wells, where ²²²Rn correlated with ²³⁸uranium (0.57). In addition, ²²²Rn is weakly correlated with total lead (0.38, p = 0.005) in the NEC aquifer (Fig. 7d). ²²²Rn and ²¹⁰Pb were also correlated (0.74) in the Finnish study. Where ²²²Rn in ground water is high it is possible that some lead associated with ²²²Rn decay remains dissolved (Focazio, 2001).

Figure 7a-d. Gross alpha is related to (a) uranium and (b) ²²²Rn. ²²²Rn is related to (c) uranium and (d) lead in New England Crystalline aquifers.

Columbia Plateau Glacial aquifers

The Columbia Plateau Glacial aquifers (CPG) are affected by agricultural irrigation (Embrey, 1988). TDS concentrations exceeding 500 mg/L (Fig. 8d) indicate irrigation effects (Whitehead, 2000). Uranium correlates with bicarbonate (0.77), field specific conductance (0.71), sodium (0.70), TDS (0.68), acid neutralizing capacity (0.68), sulfate (0.51), and pH (-0.48) (Fig. 8a-c). These suggest that complexation with bicarbonate and sulfate ions (Hem, 1985) is a control on uranium solubility.




Figure 8a-d. Uranium is related to (a) bicarbonate, (b) TDS, and (c) pH; Nitrate is related to TDS (d) in the Columbia Plateau Glacial aquifer.

Uranium and ²²²Rn in principal aquifers

Radionuclide concentrations differ by Principal Aquifer (Table 1). The median uranium concentration in the three Principal Aquifers in this study are below 1 microgram per liter (µg/L), but the 75th and 90th percentiles vary. Viewed this way, uranium is highest in New England Crystalline aquifers (Table 1) and lowest in the Cambrian – Ordovician aquifer. The median ²²²Rn concentration is near the USEPA proposed MCL of 300 Pico curies per liter (pCi/L) in the glacial and Cambrian – Ordovician aquifers but is 7 times higher (2,144 pCi/L) in New England Crystalline aquifers where 25% of wells sampled exceed the proposed USEPA alternative MCL (AMCL) of 4,000 pCi/L (Table 1).

Table 1. Summary of uranium and ²²² Rn by Principal Aquifer in the glaciated northern United States									
Principal Aquifer	N	Maximum	75 th	Median	25 th	10 th	Minimum	Uranium, µg/L	²²² Rn, pCi/L
New England Crystalline	113	428.9	12.8	2.9	<1	<1	<1	<1	<80
Glacial	874	162.3	7.6	2.1	<1	<1	<1	<1	<80
Cambrian-Ordovician	73	5	3	1	<1	<1	<1	<1	<80
²²²Rn, pCi/L									
New England Crystalline	107	215,210	12,000	4,899	2,144	1,200	573	<80	<80
Glacial	609	8,056	888	523	313	208	150	<80	<80
Cambrian – Ordovician	70	2,700	920	616	370	230	181	<80	<80

Sources of uranium and ²²²Rn

Variation of uranium and ²²²Rn in ground water depends in part on the source materials of the aquifers. The source of glacial materials depends on factors that include 1) the direction of glacial ice lobe movement; 2) existence of older glacial deposits; and 3) the underlying bedrock geology. Ice that flowed into the upper Midwest, United States from Canada came from various directions and carried materials from along its flow path. This material is a mix of existing glacial deposits (which may have originated elsewhere) and materials from the local geology, creating a composite mixture that can affect contaminant concentrations, such as uranium and ²²²Rn.