

Water Resources Research Center Annual Technical Report FY 2001

Introduction

Water Resources Research Center

Annual Technical Report

FY 2001 Introduction

WATER PROBLEMS AND ISSUES OF MISSOURI

The water problems and issues in the State of Missouri can be separated into three general areas: 1) water quality, 2) water quantity, and 3) water policy. Each of Missouri's specific problems usually requires knowledge in these three areas.

Water Quality: News media attention to the occurrence of pesticides in drinking water in the Midwest has raised a serious public concern over the quality of Missouri drinking water and how it can be protected. With the large agricultural activity in the state, non-point source pollution is of major interest. Because of several hazardous waste super-fund sites, hazardous waste is still of a concern to the public. The Center's research has been to evaluate the quality of current waste sources and improve the methods to protect them. Areas of research for the past ten years have included (but are not limited to): erosion, non-point pollution, reclamation of strip mine areas, hazardous waste disposal, acid precipitation, anthropogenic effects on aquatic ecosystems and wetlands.

Water Quantity: Missouri has a history of either inadequate amounts of rainfall, or spring floods. Because of the 1987-1989 drought years, and the flood of 93 and 95, water quantity has become a major topic of concern. Research is needed to better understand droughts and flood conditions. **Water Policy:** Policies and programs need to be formulated that will ensure continued availability of water, as new demands are placed on Missouri water. The social and economic costs may no longer be held at acceptable levels if water becomes a major issue in cities and rural areas. Past droughts and the possible lowering of the Missouri River have raised serious questions over states' rights to water and priority uses. Research areas in this program have included drought planning, legal aspects, perception and values, economic analysis, recreation, land/water use policy and legislation, and long-term effects of policy decision.

COOPERATIVE ARRANGEMENTS

The following individuals have participated in the selection and development of our 2001 research program. They have been active advisory committee members, participating in research meetings and assisting with their expertise in the area of water research, and at Center research meetings. Five proposals were submitted for regional competition. Of those five, three were funded; one of which was a continuation from last year.

UNIVERSITY OF MISSOURI FACULTY ADVISORY COMMITTEE

Craig Adams University of Missouri-Columbia I-29 Agriculture Building Columbia, MO Mike Chippendale University of Missouri - Columbia 1-29 Agriculture Building Columbia, MO 65211 Dr. John Gardner Associate Dean of Research & Outreach College of Agriculture, Food & Natural Resources 2-44 Ag Building Columbia, MO Dr. Patrick Osbourne University of Missouri - St. Louis 224 Research Building St. Louis, MO 63121 Dr. Jerry Richardson University of Missouri-Kansas City 00126 RHFH 574 Kansas City, MO

STATE OF MISSOURI ADVISORY COMMITTEE MEMBERS

Dr. John Madras Department of Natural Resources Water and Pollution Control PO Box 176 Jefferson City, MO 65102 Dr. Steve Mellis Agricultural Engineering Extension 227 Agricultural Engineering University of Missouri-Columbia Columbia, MO 65201 Dr. Russell Rhodes Southwest Missouri State University 901 South National Springfield, MO 65802 U.S. Geological Survey 1400 Independence Road Rolla, MO 65401 Steve McIntosh Water Resources Program Department of Natural Resources PO Box 176 Jefferson City, MO Hamed Mubarak 205 Jefferson Street Jefferson City, MO 65101 City, MO 65102 Jim Czarneszki Fisheries & Wildlife Department of Conservation 1110 South College Ave. Columbia, MO 65201 Becky Shannon MO Department of Natural Resources PO Box 176 Jefferson City, MO 65102

Research Program

PROGRAM GOALS AND PRIORITIES

The Missouri Water Resources Research Centers goals are: 1) establish active research programs to aid in understanding and solving Missouri's and the nation's water problems; 2) provide educational opportunities in research for students with an interest in water resources and related fields; and 3) be actively dedicated to the dissemination of water related information, using all aspects of the media.

With these goals, the Center is able to mobilize the best faculty expertise in the state to examine specific water resources problems. The Center is familiar with research needs and activities, and its goals are to help researchers avoid duplicate efforts and to serve as a link between the research community and potential users of research results - such as industries, planning commissions, and state agencies.

Because of Missouri's economy revolves around its water resources, the director and principal investigators have worked closely with the state in addressing their problems by providing research data which are necessary in order to solve present and future water problems. Each of the research projects forwarded for regional competition has undergone a thorough evaluation process by the Water Centers Advisory Committee to determine its importance in solving Missouri's and the nation's water problems.

Research Program

Research Program

PROGRAM GOALS AND PRIORITIES

The Missouri Water Resources Research Centers goals are: 1) establish active research programs to aid in understanding and solving Missouri and the nations water problems; 2) provide educational opportunities in research for students with an interest in water resources and related fields; and 3) be actively dedicated to the dissemination of water related information, using all aspects of the media.

With these goals, the Center is able to mobilize the best faculty expertise in the state to examine specific water resources problems. The Center is familiar with research needs and activities, and its goals are to help researchers avoid duplicate efforts and to serve as a link between the research community and potential users of research results - such as industries, planning commissions, and state agencies

Because of Missouri's economy revolves around its water resources, the director and principal investigators have worked closely with the state in addressing their problems by providing research data which are necessary in order to solve present and future water problems. Each of the research projects forwarded for regional competition has undergone a thorough evaluation process by the Water Centers Advisory Committee to determine its importance in solving Missouri and the nations water problems.

Development of a Simple Combustion Process for Disposal of Waste from Livestock Operations

Basic Information

Title:	Development of a Simple Combustion Process for Disposal of Waste from Livestock Operations
Project Number:	2000MO3B
Start Date:	3/1/1999
End Date:	8/31/2001
Funding Source:	104B
Congressional District:	Missouri 9th
Research Category:	Water Quality
Focus Category:	Treatment, Non Point Pollution, Agriculture
Descriptors:	Animal Waste, Wastewater, Agriculture, Waste Disposal, Pollution Control, Treatment
Principal Investigators:	Virgil Flanigan, Shubhender Kapila

Publication

Progress Report (01)

Development of a Simple Combustion Process for Disposal of Waste from Livestock Operations

Project No: **USDI-1434-HQ-96GR-02680-04**
Development of a Simple Combustion Process for Disposal of
Waste from Livestock Operations
PI: Virgil Flanigan and Shubhen Kapila

Project Period: March 1, 1999 – February 28, 2001

Description:

The project was initiated in March 1999. The first four months were spent on design and fabrication of the combustor. The combustor has now been built and installed at the Center for Environmental Science and Technology at the University of Missouri-Rolla.

The optimization and evaluation of the burner configuration will be carried out in the next three months. This will be followed up with trial degradation of simulated waste. These evaluations will include chemical and biological measurements to ascertain degradation efficiencies of odiferous chemicals, pathogens, pesticides, residual antibiotics and metabolites. The analytical methodologies required for the measurements are being validated.

We are planning to carry out a trial with farm waste streams during the summer and fall 2000. A complete project report will be submitted by March 2001.

Identifying Reference Stream Reaches for Comprehensive bioassessment

Basic Information

Title:	Identifying Reference Stream Reaches for Comprehensive bioassessment
Project Number:	2001MO2901B
Start Date:	7/1/2001
End Date:	6/30/2002
Funding Source:	104B
Congressional District:	9th
Research Category:	Not Applicable
Focus Category:	Water Quality, Non Point Pollution, Nutrients
Descriptors:	bioassessment, water quality,nutrients
Principal Investigators:	Charles F. Rabeni, Charles F. Rabeni

Publication

Summary Report of Final Project

Identifying reference stream reaches for comprehensive bioassessment.

Charles F. Rabeni
Missouri Cooperative Fish and Wildlife Research Unit
Department of Fisheries and Wildlife Sciences
University of Missouri
Columbia Missouri 65211

Research was conducted to validate the Missouri Resource Assessment Partnership's (MoRAP) newly designed hierarchical classification scheme using biotic assemblages. The Aquatic Ecological System (AES) level of the scheme was evaluated using crayfish assemblages. Six species of crayfish were sampled from four macrohabitats: runs, backwaters, riffles, vegetation plots, from nine streams in three AES units within the Meramec Ecological Drainage Unit (EDU). Crayfish assemblages were compared both within and between AES's for all four macrohabitat types.

The results showed that there were differences in crayfish assemblages between the three AES Types, just as predicted by the model. Assemblages of crayfishes varied between the three AES Types for any given macrohabitat, yet the assemblages, within an AES Type, were relatively similar for riffles and runs as well as backwaters and vegetation plots. There was more variation in assemblages between AES types than within, suggesting that the Aquatic Ecological System level of the hierarchy was appropriate for classifying differences in some aquatic biota. This research indicates a good potential for the MoRAP system in improving conservation and management programs at the regional level. Subsequent evaluation of the hierarchical system using the benthic invertebrate community is currently underway.

Identification and Biological Screening of Endocrine Disruptors in Effluents from Missouri Sewage Treatment Plants

Basic Information

Title:	Identification and Biological Screening of Endocrine Disruptors in Effluents from Missouri Sewage Treatment Plants
Project Number:	2001MO3002B
Start Date:	3/1/2001
End Date:	2/28/2002
Funding Source:	104B
Congressional District:	8th
Research Category:	Not Applicable
Focus Category:	Water Quality, None, None
Descriptors:	Water Quality, Sewage,Endocrine Disruptors
Principal Investigators:	Yue-wern Huang, Paul K.S. Nam

Publication

Identification and Biological Screening of Endocrine Disruptors in Effluents from Missouri Sewage Treatment Plants

PI: Yue-wern Huang, Department of Biological Sciences and Center for Environmental Sciences and Technology, University of Missouri-Rolla

Co-PI: Paul Nam, Center for Environmental Sciences and Technology, University of Missouri-Rolla

Funding Agency:
U.S. Geological Survey
Missouri Water Resources and Research Center

Grant Number: 01HQGR0089

Contact Person:
Yue-wern Huang, Ph.D.
Assistant Professor of Environmental Toxicology
Department of Biological Sciences
University of Missouri-Rolla
105 Schrenk Hall
1870 Miner Circle
Rolla, MO 65409

Tel: 573-341-6589
Fax: 573-341-4821
E-Mail: huangy@umr.edu

TABLE OF CONTENTS

	<u>Page</u>
Cover.....	1
Table of Contents.....	2
List of Figures.....	3
List of Tables.....	4
Abstract.....	5
Introduction.....	5
Methods and Materials.....	6
Results and Discussions.....	9
References.....	12
Appendix.....	22

LIST OF TABLES

<u>Table No.</u>	<u>Page</u>
1. Quality Control for Chemical Analysis of the STP Effluents.....	13
2. Occurrence and Levels of Chemicals in Missouri Sewage Treatment Plants on June 27-28 2001.....	14
3. Occurrence and Levels of Chemicals in Missouri Sewage Treatment Plants on August 8-9 2001.....	15
4. Occurrence and Levels of Chemicals in Missouri Sewage Treatment Plants on October 18-19 2001.....	16
5. Estrogenic Response of MCF7 Cells to Environmental Chemicals.....	17

LIST OF FIGURES

<u>Figure No.</u>		<u>Page</u>
1.	Representative concentration-response curves of chemicals with strong estrogenicity	18
2.	Representative concentration-response curves of chemicals with weak estrogenicity	19
3.	A representative concentration-response curve of permethrin, a weak estrogen, on the proliferation of MCF7 cells.....	20
4.	Representative concentration-response curves of the effluent extracts collected on Oct. 18 - 19, 2002.	21

ABSTRACT

Effluent samples were taken from four Missouri sewage treatment plants (STP). Each effluent sample was subject to a solid phase extraction (SPE) and then analyzed for endogenous estrogens, synthetic estrogens, pesticides, herbicides, industrial chemicals, and α -zearalenol (a mycotoxin) using LC/MS and GC/MS. An aliquot of each sample from the SPE were reconstituted in DMSO, which were further diluted with medium for testing their ability to induce MCF-7 cell proliferation. The proliferation data from each reconstituted water extract were compared with 17β -estradiol (E2), a standard curve. Chemicals that were detected and/or have been reported to be estrogenic were also tested individually to calculate EC50 values, relative proliferative potency (RPP), and relative proliferative effect (RPE) in stimulating MCF7 cell proliferation (Figures 1 – 3).

The chemical detection limits are between 0.1 ng/L and 0.9 ng/L, whereas the recovery of the chemicals are between 51.7% and 93.3% (Table 1). Among the endogenous and synthetic estrogens, only 17α -ethinylestradiol was found once in the Columbia STP at 2.9 ng/L water (Tables 2 – 4). In general, industrial chemicals such as 4-octylphenol, nonylphenols, 4-tert-octylphenol, dibutyl phthalate, butyl benzyl phthalate, and bisphenol A are more prevalent in the effluents compared with the other types of chemicals (Tables 2 – 4). However, these industrial chemicals are less estrogenic than endogenous and synthetic estrogens (Table 5). Figure 1 shows that the effluent extracts from the Little Blue Valley, Kansas City, and Columbia STPs are capable of stimulating MCF7 cell proliferation. The induction can be inhibited by antiestrogen (i.e. tamoxifen) co-treatments (data not shown), indicating that the cell proliferation is mediated via an estrogen receptor (ER) signal transduction pathway. The estrogenic effects might be caused by industrial chemicals detected and/or other chemicals that were not identified in our chemical analysis. Whether the effluents would have imposed estrogenic effects on organisms inhabiting the STP downstream ecosystems remains to be investigated.

INTRODUCTION

Estrogen, an endogenous sex steroid hormone, plays a major role in secondary sex organ development, behavior, fertility, and reproductive capacity. Many environmental pollutants such as herbicides, pesticides, phytochemicals, industrial wastes, and pharmaceuticals have been reported to possess estrogen-like activity (1). It has been shown that exposure to these xenoestrogens causes abnormalities in reproductive organs and malformations in wildlife (1-3). For instance, studies in England (4-6) and the USA (7) have shown that male fish held in treated sewage effluents exhibit increased levels of vitellogenin, an indication of exposure to estrogenic chemicals.

Sewage treatment plants (STPs) receive influent from domestic, municipal, and industrial sewage systems. The estrogenic chemicals in the influent may include endogenous 17β -estradiol (E2) and its metabolites (i.e., estrone, estriol), synthetic estrogens such as 17α -ethinylestradiol, personal care products, and industrial discharges such as alkylphenols. Pyrethroid insecticides and herbicides in domestic use may also end up in the sewage treatment plants. To date, the existence of estrogenic chemicals and their levels in the output of the metropolitan sewage treatment plants in Missouri are still

unsubstantiated. The potential impact of these chemicals on the quality of surface and ground waters, public health, and aquatic ecosystems remains to be elucidated. In this study we planned 1) to identify the estrogenic chemicals that may be present and determine their levels in Missouri STP effluents; and 2) to evaluate the total estrogenicity induced by effluent extracts from different STPs using the MCF7 cell proliferation test.

Effluents were collected from four sewage treatment plants in Missouri to identify and determine the levels of estrogenic chemicals. Sixteen chemicals that are endogenous estrogens, synthetic estrogens, herbicides, pesticides, and industrial chemicals were analyzed using liquid chromatography/mass spectrometry (LC/MS) and gas chromatography/mass spectrometry (GC/MS). The water extracts were subject to the MCF7 cell proliferation test that has at least two merits (8). First, the reported detection limit of 10 pg/ml E2 makes MCF7 cell proliferation assay one of the most sensitive *in vitro* assays for assessing the estrogenicity of xenoestrogens. Second, there have been few reported cases of false positive results using the MCF7 E-Screen.

METHODS AND MATERIALS

17 α -Ethinylestradiol (98%), estrone (99.3%), and estriol (99%) were purchased from ICN Biomedicals (Aurora, CA, USA). 17 β -Estradiol (E2, \geq 98%), β -estradiol-3-benzoate (EB, 98%), diethylstilbestrol (DES, \geq 99%), and dimethyl sulfoxide (DMSO, >99.5%) were purchased from Sigma (St. Louis, MO, USA). α -Zearalenol (97%), dibutyl phthalate (DBP, >99%), bisphenol A (Bp A, >99%), and 4-tert-octylphenol (97%) were purchased from Aldrich (Milwaukee, WI, USA). Butyl benzyl phthalate (BBP, 98%), atrazine (98%), simazine (98%), fenvalerate (99%, cis), bioallethrin (d-trans-allethrin, cis: trans = 2 : 96), and permethrin (cis : trans = 20:78) were purchased from ChemService (West Chester, PA, USA). SPE cartridges (Part # 188-1360) containing 1g of C₁₈ adsorbent each were purchased from J&W Scientific (Folsom, CA). The MCF7 breast cancer cell line was purchased from the Karmanos Cancer Institute (Detroit, MI). DMEM F12 medium, calf serum, and penicillin/streptomycin were purchased from Life Technologies (Grand Island, NY, USA). All other chemicals and reagents were of the highest quality available from commercial sources.

Sewage Treatment Plants and Effluent Sample Collection

Approximately 4 - 8 liters of representative 24-h composite samples of effluents from municipal sewage treatment plants in St. Louis (Bissell Plant), Columbia, Kansas City in Missouri (Blue River Plant), and Independence (Little Blue Valley Plant) were taken with an automatic, time-proportioned sampling device (Isco 3710, Lincoln, NE, USA) in June, August, and October 2001. All water samples were placed in pre-cleaned glass bottles and stored in the freezer (-20°C) until testing. These four plants have mechanical purification, active sludge treatment, biological nitrate removal, and settlement tanks as major cleaning processes.

Solid Phase Extraction

Within 2 - 3 days after collection, organic materials were recovered from the water sample by a solid phase extraction (SPE) using octadecylsilane (C₁₈) coated supports. Five mL of methanol was added to 1 liter of water sample. The water sample

was then filtered through a vacuum filtration apparatus with a glass microfiber filter (Part # 1823 047, Whatman International, Maidstone, England). The filtered water sample was pulled through a conditioned SPE cartridge at a flow rate of 10-15 mL/min by adjusting the vacuum. The SPE cartridge was washed with 6mL of deionized water, and the SPE adsorbents were dried by pulling air for a while and storing in a desiccator. The dried SPE cartridge was connected to a disposable pipette filled with anhydrous sodium sulfate. The organic materials in the SPE cartridge were eluted into an empty graduated tube by passing 3 mL of acetone twice. The extract was concentrated to 1mL under gentle stream of nitrogen. Half of the extract was transferred to a tube for further chemical analysis. Fifty μ L of DMSO was added to the remaining extract, and acetone was completely removed under a gentle stream of nitrogen. This fraction of water was stored at -70°C for the subsequent cell proliferation assay.

Chemical Analysis

The extracted organics were analyzed for the presence of estrogenic chemicals such as steroidal estrogens, pesticides, herbicides, and industrial chemicals using LC/MS and GC/MS techniques.

1. Silica Gel Fractionation

Prior to LC/MS and GC/MS analyses, the SPE samples were fractionated with a silica gel column which contained 1 gram of silica gel 60 sorbent (70-230 mesh, Fisher Scientific, Pittsburgh, PA) deactivated with 1.5% water. Before adding the sample, the silica gel column was rinsed with 10 mL of hexane/acetone (60:40) mixture. After transferring SPE sample to the column, the analytes were eluted with 10 mL of hexane/acetone (60:40) mixture.

2. LC/MS Analysis

A liquid chromatography/mass spectrometry (LC/MS) analysis of the analytes was performed with Hitachi M-8000 LC/MS system. Compounds of interest were separated on a reversed phase LC column (Xper-Chrom, 4.6 mm X 25 cm, P.J. Cobert Associates, Inc., St. Louis, MO) packed with C_{18} coated 5 μ supports. A 35-minute gradient program of 100% to 0% water using two mobile phases (water and acetonitrile with 1% acetic acid additive) was used for the separation of the analytes. The effluent from the column was split and sent to both MS and diode array detectors. The diode array detector wavelength was set at 220 nm. The ion trap mass spectrometer is operated in positive and negative modes with an electrospray ionization source.

3. GC/MS Analysis

A gas chromatography/mass spectrometry (GC/MS) analysis of the SPE sample was performed using a capillary column (30m x 0.25mm i.d.) coated with 5% phenyl – 95% methyl polysiloxane stationary phase (DB-5, J&W Scientific, Folsom, CA). A temperature program was started from 120 $^{\circ}\text{C}$ and increased to 190 $^{\circ}\text{C}$ at 10 $^{\circ}\text{C}/\text{min}$ rate and further increased to 300 $^{\circ}\text{C}$ at 5 $^{\circ}\text{C}/\text{min}$. A Hewlett-Packard 5971 GC/MS system was operated in both scan (45-550 amu) and selected ion monitoring modes for identification and quantitation of the analytes.

The quality of all analytical procedures was controlled by inclusion of validation samples with reference spikes, replicates, and blanks. The blank was demineralized water to check whether the clean-up step introduce estrogenic contaminants into the water samples. The blank was also used in the cell proliferation to test its estrogenicity. Table 1 showed that the detection limits ranged from 0.1 ng/L to 0.9 ng/L, whereas the recovery percentage ranged from 51.7% to 93.3%.

Subsequently, the organic materials recovered from the solid phase extraction were assayed using the MCF7 cell proliferation test.

Preparation of Water Extract for the Cell Proliferation Test

4.95 ml steroid-free experimental medium were added to each water extract, vortexed, and the solution was sterile filtered through a 0.22 μm membrane. These stock solutions containing 1% (v/v) DMSO further diluted to 10- to 5,000 fold (0.05 – 25 L final volume) with steroid-free experimental medium using sterile 50 ml polypropylene tubes. The maximum DMSO or ethanol concentrations in the final medium were kept at 0.1% for every dilution, a concentration which did not affect cell yield (9).

MCF7 Cell Proliferation Assay

This assay used the human ER-positive MCF7 breast cancer cell line (Karmanos Cancer Institute, Detroit, MI) to quantitatively determine the total estrogenic activity of the above STP water extracts. The protocol in this present study was adapted from other publications (8-10) and described as follows.

Cells were maintained in DMEM F12 medium supplemented with 10% calf serum and 100 IU/ml penicillin, and grown at 37°C in a 4% CO₂ humidified environment. Cells were inoculated into 12-well plates at a density of 6,000 cells per well and allowed to attach for 24 h. After 24 hours, the medium was aspirated and replaced with 1 mL experimental medium treated with dextran-coated charcoal to remove all steroids. Single chemicals were prepared in either DMSO or ethanol, depending on their solubility in these two solvents. Single chemicals and STP effluents were tested in a series dilution containing 6 to 8 concentrations, with each dilution tested in triplicate per assay. Three wells containing appropriate solvent but without test chemicals were used as negative control. 17 β -Estradiol (in either DMSO or ethanol) between 10⁻¹⁴ M and 10⁻⁸ M were used as positive control in each assay. Concentrations ranging from 10⁻¹³ M and 10⁻⁴ M were used to test individual chemicals that have been reported to be estrogenic and or commonly present in STP effluents or in environmental samples. Additionally, a fixed ratio dilution of each effluent sample was tested together with 5 μM of the antiestrogen tamoxifen to validate that the induction of cell proliferation was mediated by an ER-mediated signal transduction pathway. Each single chemical or effluent was tested at least three times.

Six days after exposure, the assay was terminated during the late exponential phase of proliferation by determination of the cell numbers in each well using the sulforhodamine B assay (10, 11). The experimental medium was discarded, the 12-well plates were washed with 500 μl cold phosphate buffered saline per well, and then fixed for at least 40 min with 200 μl cold 10% (w/v) trichloroacetic acid (TCA). After TCA was discarded, the cells were washed three times with tap water and dried completely

under a hood at the room temperature. Staining of the cells was performed by adding 250 μ l solution of 0.2% sulforhodamine B in a 1% acetic acid to each well. After 20 minutes, the staining solution was discarded and the cell were washed several times with 1 % acetic acid until the washing solution was colorless. After complete drying, the dye was dissolved in 300 μ l cold 10 mM Tris buffer (pH 10.5) per well and extinction at 550 nm (reference 630 nm) was measured in triplicate per well with a microplate reader (FLOURstar, BMG Labtechnologies, Durham, NC, USA) by transferring aliquots of 100 μ l into wells of 96-well plates.

Quantitative Evaluation and Statistics

Each chemical was tested at least three times. The data from the cell proliferation assay were fitted into a sigmoidal dose response equation with a variable slope (a.k.a. four-parameter logistic equation). EC50 was defined as the concentration of a chemical that induced cell proliferation half way between the baseline (i.e., bottom) and the maximal level (i.e., top). The equation was expressed as follows and the EC50 value was then derived from it.

$$Y = \text{Bottom} + \frac{(\text{Top} - \text{Bottom})}{1 + 10^{(\log \text{EC}_{50} - X) * \text{HillSlope}}}$$

We adjusted the EC50 of a chemical with the EC50 of E2 to indicate its relative proliferative potency (RPP), and it was calculated as follows.

$$\text{RPP} = \text{EC}_{50}[\text{E2}] / \text{EC}_{50}[\text{test chemical or effluent}]$$

In addition to the EC50 values and its comparison against E2, fold induction was also taken into consideration. The relative proliferative effect (RPE) indicated the relative proliferative response of a specific chemical compared with that of E2. Thus if a chemical was a full agonist, its RPE was 100. If a chemical was a partial agonist, its RPE was below 100. The RPE was determined as follows.

$$\text{RPE} = (\text{Maxi induction fold of a chemical or an effluent}) / (\text{Maxi. induction fold of E2})$$

The probit regression and calculation of EC₅₀ values was performed using GraphPad Prism software (GraphPad Software Inc., San Diego, CA).

RESULTS & DISCUSSIONS

Chemical Data

Twenty-four hour composite water effluents from the STPs in St. Louis, Columbia, Independence, and Kansas City (in Missouri) were collected on June 27-28, August 8-9, and October 18-19. The samples were subjected to LC/MS and GC/MS analyses, and the chemical data were shown in Tables 2 – 4. Endogenous and synthetic

estrogens were below detection limits, except for 17 α -ethinylestradiol which was found only once at a concentration of 2.9 ng/L water in the Columbia STP effluent (Table 4). Permethrin and bioallethrin were not found. Atrazine was detected at low single digit ng/L levels in all of the STPs, whereas simazine was found only once at 2.3 ng/L water in the Independence STP (Table 3). 4-Octylphenol and 4-tert-octylphenol were either below the detection limits or at less than 10 ng/L water. Compared with 4-octylphenol and 4-tert-octylphenol, nonylphenols were present in all of the plants during our survey, and could be as high as 226.1 ng/L water (Table 4). The levels of dibutyl phthalate were between 2.9 ng/L water and 5.1 ng/L water in the first sampling trip (Table 2), whereas its levels were higher between 3.5 ng/L water and 35.2 ng/L water in the two subsequent sampling events (Tables 3 – 4). Butyl benzyl phthalate was detected in all of the plants at the lower single digit ng/L levels in each sampling event. Bisphenol A was detected twice in the Columbia plant at 18.4 ng/L and 68.5 ng/L, and was not found in the other three plants. Overall, the Columbia plants tend to have higher concentrations of industrial chemicals. In general, industrial chemicals (i.e., 4-octylphenol, nonylphenols, 4-tert-octylphenol, dibutyl phthalate, butyl benzyl phthalate, and bisphenol A) were more prevalent and had higher levels in the effluents compared with the endogenous estrogens, synthetic estrogens, pesticides, and herbicides.

Biological Data

The estrogenic responses of MCF7 cells to single chemicals tested were shown in Table 5. The average maximal induction folds of 17 β -estradiol (E2 in EtOH), estriol, 17 α -ethinylestradiol, β -estradiol benzoate (β -EB), and diethylstilbestrol (DES) were between 5 – 6 folds, whereas estrone was at 3.91 fold (Figure 1 & Table 5). E2, estriol, 17 α -ethinylestradiol, β -EB, and DES showed similar relative proliferative effects (RPE). On the other hand, the RPE value of estrone was approximately 24% lower than those of E2 and others.

Though in general the above endogenous and synthetic estrogens possessed similar maximal induction folds and RPE values, their EC50 values and relative proliferative potencies (RPP) showed significant discrepancy ranging from 0.02 to 0.94, with 17 α -ethinylestradiol the highest and estrone the lowest (Table 5). These data indicated the importance of taking both RPE and RPP into consideration when estrogenicity was compared among chemicals. The EC50 and RPE values of mycotoxin, α -zearalenol, were similar to those of estriol. In summary, the above chemicals can be classified as strong environmental estrogens as they were able to illicit high levels of cell proliferation.

Toxicity was observed at 10⁻⁴ M in 4-tert-octylphenol, nonylphenol, and Bisphenol A. At this concentration the cells in the wells were completely lost. Due to toxicity at 10⁻⁴ M, these three chemicals showed incomplete sigmoidal growth curves. Thus their EC50 values and average maximal induction folds were calculated based on the assumption that 10⁻⁵ M of E2 stimulated the highest induction of cell proliferation (Figure 2 & Table 5). Nonylphenols induced 3.94-fold and 3.19-fold cell proliferation at 10⁻⁶ M and 10⁻⁵ M, respectively, with an EC50 value of 0.45 \pm 0.61 μ M, which was about 10⁵-fold higher than that of E2. Bisphenol A induced approximately 3.8-fold and 3.4-fold cell proliferation at 10⁻⁶ M and 10⁻⁵ M, respectively, with an EC50 value of 0.28 \pm 0.05 μ M. 4-tert-Octylphenol induced 4.2-fold and 3.6-fold cell proliferation at 10⁻⁶ M

and 10^{-5} M, respectively, with an EC50 value of 0.31 ± 0.15 M. Permethrin induced 2.97-fold cell proliferation and the EC50 value was 1.66 ± 1.46 μ M (Figure 3 & Table 5). One difference between permethrin and the above three chemicals was that the cells treated with permethrin at 10^{-4} M showed a similar growth rate as those of the control group, indicating some degree of cytotoxicity at this concentration. Overall, these four chemicals showed similar estrogenicity based on their similar RPP and RPE values, though bisphenol A had a bit higher RPE and a lower RPP. In summary, compared to E2, these four industrial chemicals were considered as weak environmental estrogens.

On the other hand, bioallethrin, fenvalerate, atrazine, simazine, and dibutyl benzyl phthalate at up to 10^{-5} M did not induce MCF7 cell proliferation (figures not shown). Toxicity was observed at 10^{-4} M of these chemicals. Because these five chemicals did not induce MCF-7 cell proliferation, they were classified as non-estrogenic chemicals.

The STP effluent samples collected on October 18-19 were solid phase extracted, and then diluted to test twice for their estrogenicity in MCF-7 cell proliferation. The St. Louis sample was lost during the sample filtration because of the problem with the vacuum system. The maximal cell proliferation induction by the Little Blue Valley and Kansas City STP effluents were approximately 2.85- and 3.05-fold, respectively, and were as high as that of E2, 3.18-fold (Figure 4). The maximal cell proliferation induction by the Columbia STP effluent was at 2.86-fold. All three effluents showed their highest induction at either 0.5-L or 1.25-L equivalent dilution. At 0.05-L equivalent dilution, the cell growth rates of these three effluents were approximately 1- to 1.5-fold of the control group, indicating possible cytotoxicity that might have reduced the growth rates. The induction could be inhibited by antiestrogen (tamoxifen) co-treatments (data not shown), indicating that the cell proliferation was mediated via an estrogen receptor-mediated signal transduction pathway. We suspected that the estrogenic effects might be caused by industrial chemicals and other chemicals that were not identified in our chemical analysis.

One technical limitation in relation to the cell proliferation test was that chemicals adsorbed in the solid phase of the effluent might be discounted. Moreover, though the MCF-7 cell proliferation by the effluents was observed, future studies may investigate whether the effluents would have imposed adverse ecological effects in the downstream ecosystems due to a significant dilution in the Missouri River.

Reference:

1. Colborn T, vom Saal FS, Soto AM. 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect* 101:378-384.
2. Fry DM, Toone CK, Speich SM, Peard RJ. 1987. Sex ratio skew and breeding patterns of gulls; demographic and toxicological considerations. *Studies in Avian Biology* 10:26-43.
3. Guillette LJ, Jr., Gross T, S., Masson GR, Matter JM, Percival HF, Woodward AR. 1994. Developmental abnormalities of the gonad and abnormal sex hormone concentrations in juvenile alligators from contaminated and control lakes in Florida. *Environ Health Perspect* 102:680-688.
4. Harries JE, et al. 1996. A survey of estrogenic activity in United Kingdom inland waters. *Environ Toxicol Chem* 15:1993-2002.
5. Harries JE, Janbakhsh A, Jobling S, Matthiessen P, Sumpter JP, Tyler CR. 1999. Estrogenic potency of effluent from two sewage treatment works in the United Kingdom. *Environ Toxicol Chem* 18:932-937.
6. Harries JE, Sheahan DA, Jobling S, Matthiessen P, Neall P, Sumpter JP, Tyler T, Zaman N. 1997. Estrogenic activity in five United Kingdom rivers detected by measurement of vitellogenesis in caged male trout. *Environ Health Perspect* 16:534-542.
7. Folmar LC, Denslow ND, Rao V, Chow M, Crain DA, Enblom J, Marcino J, Guillette LJ. 1996. Vitellogenin induction and reduced serum testosterone concentrations in feral male carp (*Cyprinus carpio*) captured near a major metropolitan sewage treatment plant. *Environ Health Perspect* 104:1096-1101.
8. Soto AM, Lin T-M, Justicia H, Silvia RM, Sonnenschein C. 1992. An "in culture" bioassay to assess the estrogenicity of xenobiotics (E-screen). *Advanced Modern Environmental Toxicology* 21:295-309.
9. Soto AM, Sonnenschein C, Chung HL, Fernandez MF, Olea N, Serrano FO. 1995. The E-screen assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ Health Perspect* 103(Suppl 7):113-122.
10. Körner W, Hanf V, Schuller W, Kempter C, Metzger J, Hagenmaier H. 1999. Development of a sensitive E-screen assay for quantitative analysis of estrogenic activity in municipal sewage plant effluents. *Sci Total Environ* 225:33-48.
11. Skehan P, Storeng R, Scudiero D, Monks A, MaMahon J, Vistica D, Warren JT, Bokesch H, Kenney S, Boyd MR. 1990. New colorimetric cytotoxicity assay for anticancer-drug screening. *Journal of National Cancer Institute* 82:1107-1112.

Table 1. Quality control for the chemical analysis of the STP effluents. Detection limit was determined based on 8-liter water samples. Concentrations were expressed as ng/L. R. Blank = Reagent Blank; R. Spike = Reagent Spike; DL = detection limit.

Chemical	Det. Limit	Mean R. Blank	Mean R. Spike % Recovery
17 β -Estradiol	0.5	< DL	66.7
Estrone	0.5	< DL	70.8
Estriol	0.9	< DL	51.7
17 α -Ethinylestradiol	0.5	< DL	79.2
β -Estradiol Benzoate	0.4	< DL	80.8
Diethylstilbestrol	0.8	< DL	72.5
Permethrin	0.5	< DL	70.0
Bioallethrin	0.3	< DL	84.2
Fenvalerate*	---	---	---
Atrazine	0.2	< DL	80.8
Simazine	0.3	< DL	77.5
4-Octylphenol	0.1	< DL	82.5
Nonylphenols	0.5	< DL	82.5
4-tert-Octylphenol	0.2	< DL	82.5
Dibutyl phthalate	0.1	1.2	80.8
Butyl benzyl phthalate	0.1	0.8	93.3
Bisphenol A	0.7	< DL	81.7
α -Zearalenol*	---	---	---

* Not analyzed, only tested for cell proliferation.

Table 2. Occurrence and levels of chemicals in Missouri sewage treatment plants on June 27-28 2001. Concentrations were expressed as ng/L. DL = detection limit.

Chemical	St. Louis	Columbia	Independence	Kansas City
17 β -Estradiol	< DL	< DL	< DL	< DL
Estrone	< DL	< DL	< DL	< DL
Estriol	< DL	< DL	< DL	< DL
17 α -Ethinylestradiol	< DL	< DL	< DL	< DL
β -Estradiol benzoate	< DL	< DL	< DL	< DL
Diethylstilbestrol	< DL	< DL	< DL	< DL
Permethrin	< DL	< DL	< DL	< DL
Bioallethrin	< DL	< DL	< DL	< DL
Fenvalerate*	---	---	---	---
Atrazine	2.9	< DL	< DL	< DL
Simazine	< DL	< DL	< DL	< DL
4-Octylphenol	< DL	< DL	1.8	2.1
Nonylphenols	4.5	5.1	18.1	43.3
4-tert-Octylphenol	< DL	< DL	< DL	< DL
Dibutyl phthalate	2.9	3.5	3.5	5.1
Butyl benzyl phthalate	1.4	1.3	3.5	3.6
Bisphenol A	< DL	< DL	< DL	< DL
α -Zearalenol*	---	---	---	---

* Not analyzed, only tested for cell proliferation.

Table 3. Occurrence and levels of chemicals in Missouri sewage treatment plants on August 8-9 2001. Concentrations were expressed as ng/L. DL = detection limit.

Chemical	St. Louis	Columbia	Independence	Kansas City
17 β -Estradiol	< DL	< DL	< DL	< DL
Estrone	< DL	< DL	< DL	< DL
Estriol	< DL	< DL	< DL	< DL
17 α -Ethinylestradiol	< DL	< DL	< DL	< DL
β -Estradiol Benzoate	< DL	< DL	< DL	< DL
Diethylstilbestrol	< DL	< DL	< DL	< DL
Permethrin	< DL	< DL	< DL	< DL
Bioallethrin	< DL	< DL	< DL	< DL
Fenvalerate*	---	---	---	---
Atrazine	4.3	< DL	1.7	4.2
Simazine	< DL	< DL	2.3	< DL
4-Octylphenol	<DL	4.8	<DL	2.0
Nonylphenols	6.5	139.5	2.9	43.6
4-tert-Octylphenol	< DL	3.2	<DL	< DL
Dibutyl phthalate	7.5	35.2	3.9	7.0
Butyl benzyl phthalate	3.0	2.0	1.9	7.8
Bisphenol A	< DL	68.5	< DL	< DL
α -Zearalenol*	---	---	---	---

* Not analyzed, only tested for cell proliferation.

Table 4. Occurrence and levels of chemicals in Missouri sewage treatment plants on October 18-19 2001. Concentrations were expressed as ng/L. DL = detection limit.

Chemical	St. Louis	Columbia	Independence	Kansas City
17 β -Estradiol	< DL	< DL	< DL	< DL
Estrone	< DL	< DL	< DL	< DL
Estriol	< DL	< DL	< DL	< DL
17 α -Ethinylestradiol	< DL	2.9	< DL	< DL
β -Estradiol Benzoate	< DL	< DL	< DL	< DL
Diethylstilbestrol	< DL	< DL	< DL	< DL
Permethrin	< DL	< DL	< DL	< DL
Bioallethrin	< DL	< DL	< DL	< DL
Fenvalerate*	---	---	---	---
Atrazine	1.9	0.6	< DL	< DL
Simazine	< DL	< DL	< DL	< DL
4-Octylphenol	0.8	6.5	4.6	1.3
Nonylphenols	13.3	226.1	101.9	30.1
4-tert-Octylphenol	1.1	3.9	3.0	1.4
Dibutyl phthalate	18.3	17.9	21.7	24.3
Butyl benzyl phthalate	5.1	3.3	4.8	5.4
Bisphenol A	< DL	18.4	< DL	< DL
α -Zearalenol*	---	---	---	---

* Not analyzed, only tested for cell proliferation.

Table 5. Estrogenic response of MCF7 cells to endogenous estrogens, synthetic estrogens, pesticides, herbicides, industrial chemicals, and one mycotoxin (α -zearalenol). The relative proliferative potency (RPP) or relative proliferative effect (RPE) values of each single chemical were calculated based on E2 in the corresponding solvent.

Chemical	EC50	RPP ¹	Ave. Max. Fold	RPE ¹
17 β -Estradiol (in EtOH)	4.16 \pm 1.84 pM	1.0	5.18 \pm 1.03	1.0
17 β -Estradiol (in DMSO)	2.16 \pm 1.68 pM	1.0	3.42 \pm 0.66	1.0
Estrone	214 \pm 82 pM	0.02 \pm 0.01	3.91 \pm 1.07	0.76 \pm 0.16
Estriol	53.5 \pm 17.0 pM	0.09 \pm 0.04	5.11 \pm 0.70	1.02 \pm 0.12
17 α -Ethinylestradiol	2.86 \pm 1.28 pM	0.94 \pm 0.23	5.38 \pm 0.56	1.08 \pm 0.15
β -Estradiol Benzoate	10.4 \pm 9.84 pM	0.71 \pm 0.69	5.14 \pm 0.88	0.93 \pm 0.22
Diethylstilbestrol	23.7 \pm 1.49 pM	0.14 \pm 0.05	5.87 \pm 1.40	1.01 \pm 0.30
Permethrin	1.66 \pm 1.46 μ M	1.22E-5 \pm 2.01E-5	2.97 \pm 1.10	0.62 \pm 0.06
Bioallethrin	N.E. ²	N.E.	N.E.	N.E.
Fenvalerate	N.E.	N.E.	N.E.	N.E.
Atrazine	N.E.	N.E.	N.E.	N.E.
Simazine	N.E.	N.E.	N.E.	N.E.
4-Octylphenol	N.T.	N.T.	N.T.	N.T.
Nonylphenols	0.45 \pm 0.61 μ M	8.85E-5 \pm 1.95E-4	3.94 \pm 0.17	0.77 \pm 0.99
4-tert-Octylphenol	0.31 \pm 0.15 μ M	2.02E-5 \pm 9.60E-6	3.93 \pm 0.66	0.81 \pm 0.15
Dibutyl benzyl phthalate	N.E.	N.E.	N.E.	N.E.
Bisphenol A	0.28 \pm 0.05 μ M	2.8E-6 \pm 2.4E-6	3.79 \pm 0.03	0.94 \pm 0.17
α -Zearalenol	59.8 \pm 73.8 pM	0.23 \pm 0.23	4.76 \pm 0.57	1.03 \pm 0.18

¹ The calculations for RPE and RPP were defined in the “*Quantitative Evaluation and Statistics*.”

² N.E. denotes not estrogenic.




Figure 1. Representative concentration-response curves of the chemicals with strong estrogenicity on the proliferation of MCF7 cells. Cells were stained with sulforhodamine B, and the cell density was determined by a spectrophotometer.




Figure 2. Representative concentration-response curves of the chemicals with weak estrogenicity on the proliferation of MCF7 cells.




Figure 3. A representative concentration-response curve of permethrin, a weak estrogen, on the proliferation of MCF7 cells.




Figure 4. Representative concentration-response curves of the effluent extracts collected on Oct. 18 - 19, 2002. Inset is the E2 in DMSO as a standard curve. The sample from St Louis STP was lost in the preparation for tissue culture experiment. The plot of the dilution volumes was log transformed. Values represent means \pm S.D. of triplicates in one single experiment.

Spatial Distribution, Geochemistry, and Sources of Phosphorus and Metals in Bottom Sediments in the James River Arm of Table Rock Lake

Basic Information

Title:	Spatial Distribution, Geochemistry, and Sources of Phosphorus and Metals in Bottom Sediments in the James River Arm of Table Rock Lake
Project Number:	2001MO3041B
Start Date:	3/1/2001
End Date:	2/28/2002
Funding Source:	
Congressional District:	7th
Research Category:	
Focus Category:	Water Quality, Non Point Pollution, Nutrients
Descriptors:	sediment, nutrients, water quality
Principal Investigators:	Robert T. Pavlowsky

Publication

Preliminary Final Report-January 27, 2003

Spatial Distribution, Geochemistry, and Sources of Phosphorus and Metals in Bottom Sediments in the James River Arm of Table Rock Lake

Dr. Robert T. Pavlowsky (PI)
Department of Geography, Geology, and Planning
Southwest Missouri State University

Federal Dollars: \$17,265

Progress Overview

All work on the project has been completed according to plan. Results of the study have been presented at two meetings: (1) White River Basin Forum, Springfield, MO, November 2001, and (2) 9th Annual Symposium on the Interactions between Sediment and Water, Banff, Canada May 2002, . The final report will take the form of a MS thesis in Resource Planning from SMSU and is currently in the final stages of preparation.

Objective 1: Watershed and Bathymetric Mapping

GIS layers for bathymetry, elevation, land use, dock locations, and geology were developed from various federal and state sources including the USGS, USACE, and MDNR. The location and elevation of the flooded James River channel and valley floor features classified from the GIS data based were augmented with GPS and depth measurements collected during sampling. The contributing drainage area and land use above tributary or cove samples was also determined for analysis of nonpoint source influence on phosphorus concentrations.

Objective 2: Bottom Sediment Characterization

The study area includes the Table Rock lake area along the 48 km length of the James River arm from Galena, Missouri to the White River. Over 200 sediment samples were collected at GPS-located sites in both the main channel of the lake and its tributary coves. Depth was determined using a hull-mounted "fish-finder" device and checked with a hand-held sonar. All sediment samples were analyzed for particle size, organic matter content, and bulk geochemistry including concentrated HCl-HNO₃ extractable phosphorus, urban metals (Cu, Hg, Pb, Zn), and substrate-forming metals (Al, Fe, Mn, and Ca).

Objective 3: Assessment of Biochemical Mobility

Nine bottom sediment samples were selected for sequential extraction procedures to determine the form of the phosphorus in the sediment. Five different P fractions were evaluated: (1) exchangeable; (2) carbonate; (3) Al/Fe hydroxides; (4) apatite; and (5) Organic or residual.

Preliminary analysis indicates that the P distribution of the sediment varies only slightly among the sites studied. The most biochemically mobile fractions (exchangeable and carbonate-bound) tend to contain between 40 and 55 % of the sediment-P in the lake and cove samples. About 20 to 30 % is found in the organic or residual fraction. There are no obvious spatial trends (downstream or depth) in the fractionation data. Generally, about half of the phosphorus in the bottom sediments of the James River arm may potentially be biologically available, depending on redox and sedimentation conditions.

Objective 4: Spatial Variability

While water-column data generally show the decay of total-P (ug/l) downstream from the James River mouth near Galena, bottom sediment P (ug/g) concentrations increase downlake suggesting that sedimentation is the major process controlling P transport to the main lake. Sediment-P concentrations are relatively variable and range from 300 to 850 ug/l along the 10 km long river-lake transition area. Over the next 38 km, P concentrations gradually increase with depth from 1,000 to 2,400 ug/g. Depending on the statistical sampling procedures and assumptions selected, sediment-P concentrations in the main stem of the arm are predictable based on depth or distance from Galena, organic matter or clay content of the sample, and concentrations of Al, Ca, Fe, and Mn in the sediment.

Relatively high concentrations of sediment-P were found in the shallow, upstream reach of the arm below McCord's Bend where algal blooms and fish kills occurred during the summer months in the past. Patterns of phosphorous distribution in the James River arm generally differed from those of metals typically released from nonpoint sources. However, urban metal levels tended to be low throughout the lake with the exception of a few high samples of Hg in the upper arm reaches and Cu near the confluence of Aunt's Creek in the lower reaches of the arm. Several samples collected from the White River portion of Table Rock Lake indicate that enriched levels of bottom sediment-P are extending out into the main lake from the James River arm.

Efforts to explain variations in tributary or cove sediment-P concentrations with land use variables including percent urban, agricultural, and forested area, dock and road density, and tributary lake and watershed area were not successful. This was due to two main factors. First, phosphorus enrichment of bottom sediments by anthropogenic sources probably occurs at levels far below the influence of natural sediment sorption processes and background source variations. Second, land use variables covaried with cove watershed size in a manner that countered source-P transport linkages in the regression analyses. For example, while larger watershed areas had more urban development due to historical settlement trends, lower slopes, and building site availability, they also were more forested and had greater potential for dilution from background sediment transport. Also, the main plume of phosphorus moving down lake was able to enter and become deposited within very small coves with low discharges; large coves had enough runoff to counter this effect or were large enough to evade detection during this study.

The results of this study suggest that phosphorus contamination problems in the main arm are mainly the result of fluvial inputs from the James River at Galena and not sources located in its direct cove drainages. However, this study did not evaluate the local effects of phosphorus

introductions from nonpoint sources in the coves on sediment-P levels. Also, this study did not resolve the effect of P treatment upgrades on Springfield's Southwest Treatment Plant, which has been the primary source of P to the arm in the past.

Objective 5: Phosphorus Budget

Published data on water-column total phosphorus concentrations in the James River arm and James River at Galena and mean annual discharge at gaging sites in southwest Missouri is used to develop a phosphorus budget for the James River arm of Table Rock Lake. This transport budget is compared with sediment-P storage to understand the role of sedimentation and sediment remobilization in the James River arm. Under average hydrologic conditions, preliminary results indicate that most of the phosphorus entering the lake at Galena and from other nonpoint sources via tributary cover inputs is removed by sedimentation or biologic uptake before reaching the main lake on the White River. The budget model indicates a slight net release to the main lake that supports previous findings of elevated bottom sediments in Table Rock Lake just downstream of the James River arm confluence. The importance of bottom sediment storage and cycling to P transport in the James River arm has not yet been investigated, but will be included in the final report in Marc Owen's master's thesis.

Information Transfer Program

Student Support

Student Support					
Category	Section 104 Base Grant	Section 104 RCGP Award	NIWR-USGS Internship	Supplemental Awards	Total
Undergraduate	3	0	0	0	0
Masters	2	0	0	0	0
Ph.D.	0	0	0	0	0
Post-Doc.	0	0	0	0	0
Total	5	0	0	0	0

Notable Awards and Achievements

Publications from Prior Projects